

### Cedarville University DigitalCommons@Cedarville

The Research and Scholarship Symposium

The 2022 Symposium

### Characterization of a Laminin-like Protein in the Kinetodesmal Fibers of Tetrahymena thermophila

Madeline Ardrey
Cedarville University

Kaylee Howell

Cedarville University

Abbigail Paterson
Cedarville University

Adin Pendell Cedarville University

Ezra Shimabenga Cedarville University

See next page for additional authors

Follow this and additional works at: https://digitalcommons.cedarville.edu/rs\_symposium

Ardrey, Madeline; Howell, Kaylee; Paterson, Abbigail; Pendell, Adin; Shimabenga, Ezra; Silveira, Nicholas; Tomic, Anna; and Kuruvilla, Heather G., "Characterization of a Laminin-like Protein in the Kinetodesmal Fibers of Tetrahymena thermophila" (2022). *The Research and Scholarship Symposium*. 7. https://digitalcommons.cedarville.edu/rs\_symposium/2022/poster\_presentations/7

This Poster is brought to you for free and open access by DigitalCommons@Cedarville, a service of the Centennial Library. It has been accepted for inclusion in The Research and Scholarship Symposium by an authorized administrator of DigitalCommons@Cedarville. For more information, please contact digitalcommons@cedarville.edu.



<b>esenters</b> adeline Ardre ina Tomic, ar	y, Kaylee Howell, Id Heather G. Kuri	Abbigail Paters uvilla	son, Adin Pend	lell, Ezra Shima	benga, Nicholas	s Silveira,

Figure 1. A. Localization of a laminin-like protein to the kinetodesmal fibers of Tetrahymena. B. Localization of centrin to the basal bodies and centrosome of Tetrahymena.





Figure 2. Colocalization of a laminin-like protein with centrin in Tetrahymena. Red: laminin, green: centrin blue: DAPI, yellow (far left) merged image showing colocalization




# A laminin-like protein localizes to kinetodesmal fibers in Tetrahymena thermophila. This protein colocalizes with centrin in immunofluorescence as well as CO-IP.

#### Abstract

Tetrahymena thermophila are free-living ciliated organisms belonging to Kingdom Protista. These organisms possess large numbers of cilia that are used for feeding as well as locomotion. The cilia are attached to microtubule organizing centers (MTOC) called basal bodies, which are rich in centrin, a calcium binding protein present in MTOCs. Because of the mechanical stress that ciliary beating puts on the plasma membrane, these cells have kinetodesmal fibers, composed of bundled coiled-coil proteins which grow from the proximal end of the basal bodies and stabilize the plasma membrane.

Laminin is a cross-linking protein found in the extracellular matrix of animals. Using an antibody against laminin-β1, we found that the antibody localized to kinetodesmal fibers which overlapped the basal bodies stained by the anti-centrin antibody. To determine whether these proteins were in the same complex, we did co-immunoprecipitation using both our anti-centrin and anti-laminin antibodies. Both antibodies precipitated centrin (a doublet at 20 kD) as well as a number of presumed laminin-like proteins between 37 and 50 kD. We also found a number of laminin homologues in the *Tetrahymena* Genome Database. Further studies will be needed to determine which of these proteins is present in the kinetodesmal fiber.

Figure 3. BLAST alignment of laminin ( $\beta$ 1) with the predicted *Tetrahymena* proteome shows many proteins, including putative transmembrane proteins, show homology with LAM\beta1.



**Figure 3A**. Results of BLAST alignment of laminin (β1) with the proteins listed in the *Tetrahymena* genome database (ciliate.org). Colored lines (top) indicate regions of alignment between the two protein sequences. High bit scores and low E values indicate a high degree of homology between the proteins being compared.

REJ domain protein

Figure 3B. Western blot shows immunoprecipate of Tetrahymena whole cell extract with anti-centrin antibody (lane 2) and anti-laminin antibody (lane 5). Both antibodies pulled down centrin (20 kDa, bottom row, lanes 2 and 5) as well as laminin-like proteins between 37 and 50 kDa (top row, lanes 2 and 5).

## Characterization of a Laminin-like Protein in the Kinetodesmal Fibers of Tetrahymena thermophila

