

Cedarville University DigitalCommons@Cedarville

The Research and Scholarship Symposium

The 2015 Symposium

Apr 1st, 11:00 AM - 2:00 PM

Effects of Smoking vs. Nicotine Replacement Therapy During Pregnancy on Childhood Health Outcomes: An Integrative Literature Review

Andrew J. McBride Cedarville University, amcbride@cedarville.edu

Kristen P. Sabo Cedarville University, ksabo@cedarville.edu

Emily D. Williams

Cedarville University, ewilliams270@cedarville.edu

Follow this and additional works at: http://digitalcommons.cedarville.edu/research_scholarship_symposium

Part of the Female Urogenital Diseases and Pregnancy Complications Commons, Maternal and Child Health Commons, and the Maternal, Child Health and Neonatal Nursing Commons

McBride, Andrew J.; Sabo, Kristen P.; and Williams, Emily D., "Effects of Smoking vs. Nicotine Replacement Therapy During Pregnancy on Childhood Health Outcomes: An Integrative Literature Review" (2015). *The Research and Scholarship Symposium*. 36. http://digitalcommons.cedarville.edu/research_scholarship_symposium/2015/poster_presentations/36

This Poster is brought to you for free and open access by DigitalCommons@Cedarville, a service of the Centennial Library. It has been accepted for inclusion in The Research and Scholarship Symposium by an authorized administrator of DigitalCommons@Cedarville. For more information, please contact digitalcommons@cedarville.edu.



Effects of Smoking vs. NRT During Pregnancy

Andrew J. McBride, Kristen P. Sabo & Emily D. Williams

Cedarville University School of Nursing

PATIENT CARE ISSUE

- Cost for neonatal complications estimated >\$350 million annually (NC DHHS, 2013).
- Approximately 10.6% of women smoke while they are pregnant (NC DHHS, 2013).
- Smoking during pregnancy can lead to premature, low birth-weight infants, or stillbirth.
- Carbon monoxide and nicotine cross the placenta and interfere with fetal oxygen supply (NIH National Institute on Drug Abuse, 2014).
- Complications due to maternal smoking include Sudden Infant Death Syndrome, growth retardation, poor lung development, obesity, and respiratory infections (CDC, 2011).

EVIDENCE-BASED PRACTICE QUESTION

Question: In pregnant women, does the use of Nicotine Replacement Therapy compared to smoking during pregnancy reduce the risk of future childhood health concerns?

- P- Pregnant women
- I- Nicotine Replacement Therapy (NRT)
- C- Smoking
- O- Reduce the risk of future childhood health concerns (respiratory complications, obesity)

REGISTERED NURSE INTERVIEW

Labor and Delivery CRN

- NRT is often used improperly by pregnant mothers.
- Fetal heart rate increases after maternal nicotine use.

NICU CRN

- Nicotine addiction in pregnant mothers can lead to nicotine addiction in neonates.
- Mothers using nicotine often give birth to neonates who are small for gestational age, low in birth weight, and have prolonged, increased heart rate.

METHODS

Key Words: pregnancy, maternal smoking, nicotine replacement therapy, nicotine, childhood health concerns, childhood obesity

Databases: Alt HealthWatch, AMED, CINAHL, Medline


Inclusion Criteria: Articles published between 2008 and 2014, articles that focused on the fetal and future childhood concerns of NRT use and smoking use during pregnancy

Exclusion Criteria: Non-human test population, year of publication prior to 2008, small sample size

SYNTHESIS OF EVIDENCE

- Smoking leads to increased chance of preterm birth & low birth weight (LBW) (CDC, 2011).
- Childhood obesity is linked to smoking during pregnancy (Behl, 2013; Bekkers, 2011; Ino, 2010, Weng, 2012).
- NRT delivers nicotine without carcinogens (Oncken, 2008; Brose, 2013).
- NRT showed positive safety outcomes but low adherence (Coleman, 2010).
- Infants born to NRT using mothers had 2x the risk of having LBW (Gaither, 2008)
- Infants born to smokers had 1.31x the risk of LBW (Gaither, 2008).
- Combination NRT increases cessation but exposes fetus to higher nicotine levels (Brose, 2013).
- Smoking cessation should be targeted at one year prior to pregnancy (Wang, 2012).

EVIDENCE-BASED PRACTICE RECOMMENDATIONS



LIMITATIONS

- Limited studies of primary evidence within the past 5 years
- Limited studies analyzing direct effects of NRT on the fetus.
- Few studies examining longevity effects of NRT on the offspring.

REFERENCES

Behl, M., Rao, D., Aagaard, K., Davidson, T. L., Levin, E. D., Slotkin, T. A., Srinivasan, S., Wallinga, D., Holloway A.C. (2013).

Evaluation of the Association between Maternal Smoking, Childhood Obesity, and Metabolic Disorders: A A National Toxicology Program Workshop Review. *Environmental Health Prospectives* 121(2), 170-180. doi:10.1289/ehp.1205404

Bekkers, M. B. M., Brunekreef, B., Smit, H. A., Kerkhof, M., Koppelman, G. H., Oldenwening, M., & Wijga, A. H., (2011). Early-Life Determinants of Total and HDL Cholesterol Concentrations in 8-Year-Old Children; The PIAMA Birth Cohort Study. *Public Library of Science*, 6(9), e25533. doi: 10.1371/journal.pone.0025533

Brose, L., McEwen, A., West, R. (2013). Association between nicotine replacement therapy use in pregnancy and smoking cessation.

**Drug and Alcohol Dependence Journal*, 132, 660-664. DOI: 10.1016/j.drugalcdep.2013.04.017

Coleman, T., Chamberlain, C., Cooper, S., & Leondardi-Bee, J. (2010). Efficacy and safety of nicotine replacement therapy for smoking cessation in pregnancy: systematic review and meta-analysis. Society for the Study of Addiction, 106, 52-61. doi:10.1111/j. 1360-0443.2010.03179.x

Cooper, S., Lewis, S., Thornton, J. G., Marlow, N., Watts, K., Britton, J., Grainge, M. J., Taggar, J., Coleman, T. (2014). The SNAP trial: a randomised placebo-controlled trial of nicotine replacement therapy in pregnancy – clinical effectiveness and safety until 2 years after delivery, with economic evaluation. *Health Technology Assessment* 18(54), vii- 128. doi: 10.3310/hta18540

Fish, L. J., Peterson, B. L., Namenek Brouwer, R. J., Lyna, P., Oncken, C. A., Swamy, G. K., Myers, E. R., Pollak, K. N. (2009).

Adherence to Nicotine Replacement Therapy Among Pregnant Smokers. *Nicotine & Tobacco Research*, 11(5), 514-518.

doi:10.1093/ntr/ntp032

Forinash, A. B., Pitlick, J. M., Clark, K., & Alstat, V. (2010). Nicotine Replacement Therapy Effect on Pregnancy Outcomes. *The Annals of Pharmacotherapy*, 44(11), 1817- 1821. doi: 10.1345/aph.1P279

Gaither, K., Brunner Huber, L., Thompson, M., Huet-Hudeson, Y. (2008). Does the use of nicotine replacement therapy during pregnancy affect pregnancy outcomes? *Maternal and Child Health Journal*, 13, 497-504. doi:10.1007/s10995-008-0361-1

N.C. infant mortality rate for 2012 released. (2013, October 3). NC DHHS Press Release:, Retrieved October 28, 2014, from http://ncdhhs.gov/pressrel/2013/2013-10-03_NC_infant_mortality_rate.htm

Ino, T. (2010). Maternal smoking during pregnancy and offspring obesity: meta-analysis. *Pediatrics International, 52,* 94-99. doi:10.1111/j.1442-200X.2009.02883.x

Oncken, C., Dornelas, E., Greene, J., Sankey, H., Glasmann, A., Feinn, R., & Kranzler, H. R. (2008). Nicotine Gum for Pregnant Smokers: A Randomized Controlled Trial. *Obstetrics & Gynecology*, 112(4), 859-867. doi: 10.1097/AOG. 0b013e318187e1ecPregnancy and smoking. (2011, February 22). Centers for Disease Control and Prevention. Retrieved October 26, 2014, from http://www.cdc.gov/healthcommunication/toolstemplates/entertainmented/tips/pregnancy smoking.html

Smoking and pregnancy—what are the risks? (n.d.). NIH National Institute on DrugAbuse. Retrieved October 26, 2014, from

http://www.drugabuse.gov/publications/research-reports/tobacco/smoking pregnancy%E2%80%94-what-are-risks
Suzuki, K., Sato, M., Ando, D., Kondo, N., Yamagata, Z. (2013). Differences in the effect of maternal smoking during pregnancy for childhood overweight before and after 5 years of age. The Journal of Obstetrics and Gynaecology Research, 39(5), 914-921. doi:10.1111/jog.12025

Titler, M. G., Kleiber, C., Steelman, V. J., Rakel, B. A., Budreau, G., Everett, L. Q., et.al. (2001). The Iowa model of evidence-based practice to promote quality care. *Critical Care Nursing Clinical of North America*, 13(4), 500.

Wang, L., Mamudu, H. M., Wu, T. (2012). The impact of maternal prenatal smoking on the development of childhood overweight in

school-aged children. *Pediatric Obesity, 8,* 178-188. doi:10.1111/j.2047-6310.2012.00103x Weng, S.F., Redsell, S.A., Swift, J.A., Yang, M., Glazebrook, C.P. (2012). Systematic review and meta-analyses of risk factors for childhood overweight identifiable during infancy. *Arch Dis Child, 97.* 1019-1026. doi: 10.1036/archdischild-2012-302263